





OMIM **PMC** Journals Book Protein Genome Structure PubMed Nucleotide Go Clear Search PubMed for Details Clipboard ✓ Limits Preview/Index History Show: 20 Send to Text Display Abstract About Entrez

Text Version

Overview

Help | FAQ Tutorial

E-Utilities

Entrez PubMed

New/Noteworthy

MeSH Database Single Citation Matcher

Clinical Queries

LinkOut Cubby

PubMed Services Journals Database

Batch Citation Matcher

1: J Biochem (Tokyo). 1996 Jul;120(1):104-10.

Related Articles, L

Purification and characterization of a marine bacterial beta-galactoside alpha 2,6-sialyltransferase from Photobacterium damsela JT0160.

Yamamoto T, Nakashizuka M, Kodama H, Kajihara Y, Terada I.

Sea Water Science Research Laboratory, Japan Tobacco Inc., Kanagawa.

A bacterial sialyltransferase, named sialyltransferase 0160, was purified from a marine bacterium that had been isolated from seawater from Sagami Bay, Kanagawa. This strain has been identified as Photobacterium damsela, and named P. damsela JT0160. Sialyltransferase 0160 was purified 688-fold to homogeneity from the crude extract of th cells with a yield of 19% using a combination of anion exchange chromatography, hydroxyapatite chromatography, gel filtration chromatography, and affinity chromatography. The purified enzyme migrated as a single band (61 kDa) on sodium dodecyl sulfate-polyacrylamide gel. This sialyltransferase was found to be a betagalactoside alpha 2,6-sialyltransferase [EC 2.4.99.1] which catalyzes the incorporation o NeuAc from CMP-NeuAc into the galactose residue of the carbohydrate chain at positio on the basis of an analysis of the enzymatic reaction products with HPLC, 1H-, 13C-NM spectroscopy, and fast atom bombardment mass spectroscopy.

PMID: 8864851 [PubMed - indexed for MEDLINE]

Order Documents NLM Catalog NLM Gateway TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central

Related Resources

h

Display Abstract	Show:	20 🔻	Sort · 🔽	Send to Text	V

Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Privacy Statement | Freedom of Information Act | Disclaimer

Oct 4 2004 14

h g cb b







PMC Book **OMIM** Journals Genome Structure PubMed Nucleotide Protein Go Clear Search PubMed for Details Preview/Index Clipboard **☑** Limits History Show: 20 Sort Send to Text Display Abstract

About Entrez

Text Version

E-Utilities

Entrez PubMed Overview Help | FAQ Tutorial New/Noteworthy

PubMed Services Journals Database MeSH Database Single Citation Matcher **Batch Citation Matcher Clinical Queries** LinkOut Cubby

Related Resources Order Documents **NLM Catalog NLM Gateway** TOXNET Consumer Health Clinical Alerts ClinicalTrials.gov PubMed Central

1: Glycobiology. 1991 Sep;1(4):357-65.

Related Articles, L

Complete nucleotide and deduced protein sequence of CMP-NeuAc: polyalpha-2,8 sialosyl sialyltransferase of Escherichia coli K1.

Weisgerber C, Hansen A, Frosch M.

Institute fur Medizinische Mikrobiologie, Medizinische Hochschule Hannover, FRG.

Poly-alpha-2,8 N-acetylneuraminic acid (polySia) is an important virulence factor in infections caused by Escherichia coli K1 and Neisseria meningitidis B. In E. coli K1 a membranous CMP-NeuAc: poly-alpha-2,8 sialosyl sialyltransferase (polysialyltransferas complex catalyses the synthesis of linear polySia chains. The complex also elongates sia oligomers that serve as exogenous acceptors. The gene encoding a polysialyltransferase E. coli has been identified by subcloning and DNA sequence analysis. The subcloned DN fragment codes for a polypeptide with a molecular mass of 47 kDa catalysing the in vitro synthesis of polySia by elongation of exogenous acceptors.

PMID: 1820197 [PubMed - indexed for MEDLINE]

Show: 20 Sort Send to Text Abstract Display



Write to the Help Desk NCBI | NLM | NIH Department of Health & Human Services Privacy Statement | Freedom of Information Act | Disclaimer

Oct 4 2004 14:

c